



# Recommendations for the Prevention of RSV Disease in Infants and Children: Policy Statement

Committee on Infectious Diseases

## RECOMMENDED POPULATIONS

The American Academy of Pediatrics (AAP) recommends respiratory syncytial virus (RSV) immunization<sup>1</sup> for:

- infants <8 months of age born during or entering their first RSV season if:
  - pregnant parent did not receive RSVpreF vaccine during this pregnancy,
  - pregnant parent's RSVpreF vaccination status is unknown, or
  - infant was born <14 days after the pregnant parent's RSVpreF vaccination
- infants and children 8 through 19 months of age at high risk of severe RSV disease and entering their second RSV season, regardless of the RSV vaccination status of the pregnant parent or the child's prior receipt of nirsevimab or clesrovimab when <8 months of age in their first RSV season. High-risk criteria include the following:
  - Children with chronic lung disease of prematurity who required medical support (chronic corticosteroid therapy, diuretic therapy, or supplemental oxygen) at any time during the 6-month period before the start of the second RSV season
  - Children with severe immunocompromise
  - Children with cystic fibrosis who have either:
    - manifestations of severe lung disease (previous hospitalization for pulmonary exacerbation in the first year of life or abnormalities on chest imaging that persist when stable), or
    - weight-for-length that is less than the 10th percentile
  - American Indian or Alaska Native children

Note that all ages refer to chronologic age, not corrected age.

## EQUITY CONSIDERATIONS

American Indian and Alaska Native Children are included in the high-risk category, because they experience significantly higher rates of severe RSV

**CONFLICT OF INTEREST DISCLOSURE:** Sean O'Leary, MD, MPH, FAAP, has no potential conflicts of interest to disclose. Claudia Espinosa, MD, MSc, FAAP, has disclosed a direct to institution financial relationship with Merck and Sanofi as a research site Principal Investigator. Dr Espinosa did not participate in making recommendations or development of this statement. Other disclosures of committee members were reviewed and determined not relevant to the work related to the statement.

**FUNDING:** No external funding.

This document is copyrighted and is property of the American Academy of Pediatrics and its Board of Directors. All authors have filed conflict of interest statements with the American Academy of Pediatrics. Any conflicts have been resolved through a process approved by the Board of Directors. The American Academy of Pediatrics has neither solicited nor accepted any commercial involvement in the development of the content of this publication.

Policy statements from the American Academy of Pediatrics benefit from expertise and resources of liaisons and internal (AAP) and external reviewers. However, policy statements from the American Academy of Pediatrics may not reflect the views of the liaisons or the organizations or government agencies that they represent.

The guidance in this statement does not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate.

All policy statements from the American Academy of Pediatrics automatically expire 5 years after publication unless reaffirmed, revised, or retired at or before that time.

<https://doi.org/10.1542/peds.2025-073923>

Copyright © 2025 by the American Academy of Pediatrics

**To cite:** American Academy of Pediatrics, Committee on Infectious Diseases. Recommendations for the Prevention of RSV Disease in Infants and Children: Policy Statement. *Pediatrics*. 2025;156(5):e2025073923

disease and hospitalization, with children living in rural and reservation communities most impacted. Evidence supports that this increased risk is associated with social drivers of health, with specific attention to running water access, transportation difficulties, household crowding, and indoor air quality.<sup>2</sup>

## APPROVED PRODUCTS

### Nirsevimab and Clesrovimab

Nirsevimab and clesrovimab<sup>3,4</sup> are considered the first-line recommended immunization products for administration to infants to protect against medically attended RSV disease. They are long-acting monoclonal antibody products administered as a single dose for administration during an infant's first RSV season. The AAP recommends any licensed RSV immunization product appropriate for age and health status and does not prefer one product over another.

- Only children 8 through 19 months who meet high-risk criteria should receive RSV immunization in their second RSV season. Nirsevimab is approved for this indication; clesrovimab is not.
- While the timing of the onset and duration of RSV season may vary, RSV immunization may be administered from October through the end of March in most of the continental United States. The timing of the onset, peak, and decline of RSV activity vary geographically, and providers may adjust timing of administration based on guidance from public health authorities (eg, Centers for Disease Control and Prevention [CDC], local health departments) or regional medical centers.
- Infants born during the RSV season should receive RSV immunization within 1 week of birth, ideally during their birth hospitalization. However, administration can occur during any visit to a health care setting, including well-child visits.
- Eligible infants with prolonged birth hospitalizations because of prematurity or other causes, who are discharged during RSV season, should receive RSV immunization shortly before or promptly after discharge from the hospital.
- Eligible infants who are <8 months of age and born outside of RSV season should receive RSV immunization shortly before or during the RSV season. Administration can occur during any visit to a health care setting, including well-child visits.
- In accordance with AAP's general best practices for immunizations, (<https://publications.aap.org/redbook>) simultaneous administration of RSV immunization with age-appropriate vaccines is recommended. In clinical trials, when RSV immunization was administered concomitantly with routine childhood vaccines, the safety and reactogenicity profile of the concomitantly administered regimen

was similar to that of the childhood vaccines administered alone. When concomitantly administered, RSV immunization is not expected to interfere with the immune response to other vaccines.

- Immunization is not needed for most infants <8 months of age whose pregnant parent received RSVpreF vaccine during the pregnancy and  $\geq 14$  days before giving birth. RSV immunization may be considered for infants born to a vaccinated pregnant parent in rare circumstances when, based on the clinical judgment of the health care provider, the potential incremental benefit of administration is warranted. These situations include, but are not limited to:
  - Infants born to pregnant people who may not mount an adequate immune response to RSV vaccination (eg, pregnant people with immunocompromising conditions)
  - Infants born to pregnant people who have medical conditions associated with reduced transplacental antibody transfer (eg, pregnant people living with HIV infection)
  - Infants who have undergone cardiopulmonary bypass (see nirsevimab FDA package insert and clesrovimab FDA package insert) or extracorporeal membrane oxygenation (ECMO), leading to loss of maternal antibodies
  - Infants with substantial increased risk for severe RSV disease (eg, hemodynamically significant congenital heart disease, intensive care admission with a requirement of oxygen at discharge)

### Palivizumab

Palivizumab is a short-acting monoclonal antibody product that is administered in monthly doses during the RSV season. Palivizumab is no longer routinely recommended for use and will be discontinued as of December 31, 2025.

### Additional Information

- RSV Immunization Administration Visual Guide ([AAP.org/rsv](https://publications.aap.org/rsv))
- RSV Immunization Frequently Asked Questions ([AAP.org/rsv](https://publications.aap.org/rsv))
- *Red Book* Chapter on Respiratory Syncytial Virus (<https://publications.aap.org/redbook>)
- RSV Immunization Information Statement (<https://www.immunize.org/vaccines/vis/iis-rsv/>)

### COMMITTEE ON INFECTIOUS DISEASES, 2025–2026

Sean T. O'Leary, MD, MPH, FAAP, Chairperson  
 James D. Campbell, MD, MS, FAAP, Vice Chairperson  
 Monica I. Ardura, DO, MSCS, FAAP  
 Kristina A. Bryant, MD, FAAP, *Red Book Online* Team Lead  
 Roberta L. DeBiasi, MD, MS, FAAP  
 Claudia Espinosa, MD, MSc, FAAP

Robert W. Frenck, Jr, MD, FAAP  
C. Mary Healy, MD  
Betsy C. Herold, MD, FAAP  
Chandy C. John, MD, MS, FAAP  
Aaron M. Milstone, MD, MHS, FAAP  
Angela L. Myers MD, MPH, FAAP  
Pia S. Pannaraj, MD, MPH, FAAP  
Adam J. Ratner, MD, MPH, FAAP, *Red Book* Associate Editor  
José R. Romero, MD, FAAP  
Matthew Zahn, MD, FAAP

## EX-OFFICIO

David W. Kimberlin, MD, FAAP, *Red Book* Editor  
Ritu Banerjee, MD, PhD, FAAP, *Red Book* Associate Editor  
Elizabeth D. Barnett, MD, FAAP, *Red Book* Associate Editor  
Ruth Lynfield, MD, FAAP, *Red Book* Associate Editor

## LIAISONS

Amina Ahmed, MD, FAAP, American Thoracic Society  
Michelle Barton, MD, Canadian Paediatric Society  
Carolyn Cleary, MD, FAAP, AAP Committee on Practice and Ambulatory Medicine  
Denée J. Moore, MD, American Academy of Family Physicians  
Jennifer L. Thompson, MD, American College of Obstetricians and Gynecologists  
Juan P. Torres, MD, PhD, Sociedad Latinoamericana de Infectología Pediátrica  
Charles R. Woods, Jr., MD, MS, FAAP, Pediatric Infectious Diseases Society

## STAFF

Sunnah Kim; Gillian Gibbs, MPH

## ACKNOWLEDGMENTS

The Committee on Infectious Diseases gratefully acknowledges Sean O'Leary, MD, MPH, FAAP, for his leadership in drafting the policy statement.

## REFERENCES

1. American Academy of Pediatrics. Respiratory syncytial virus. In: Kimberlin DW, Banerjee R, Barnett ED, Lynfield R, Sawyer MH, eds. *Red Book: 2024 Report of the Committee on Infectious Diseases*. 33rd ed. American Academy of Pediatrics; 2024:713–721.
2. Atwell JE, Hartman RM, Parker D, et al. RSV among American Indian and Alaska Native children: 2019 to 2020. *Pediatrics*. 2023;152(2):e2022060435. PubMed doi: 10.1542/peds.2022-060435
3. Zar HJ, Simoes E, Madhi S, et al. 166. A phase 2b/3 study to evaluate the efficacy and safety of an investigational respiratory syncytial virus (RSV) antibody, clesrovimab, in healthy preterm and full-term infants [abstr]. *Open Forum Infect Dis*. 2025;12(Suppl 1): ofae631.003. doi: <https://doi.org/10.1093/ofid/ofae631.003>
4. Moulia D; Maternal/Pediatric RSV Work Group. Centers for Disease Control and Prevention. Evidence to Recommendations Framework: Clesrovimab. Accessed August 8, 2025. <https://www.cdc.gov/acip/downloads/slides-2025-04-15-16/02-Moulia-maternal-peds-RSV-508.pdf>